

# Designing Feeding Systems for Robotic Milking

Presented at the Tri-State Nutrition Conference,  
April 19-20, 2011, Fort Wayne, Indiana

*Jack Rodenburg*<sup>1</sup>  
DairyLogix

**Summary** – Feed, is the primary motivation for the cow to visit the robotic milking stall. Highly motivated cows will visit voluntarily thereby decreasing the need to expend labour fetching cows, and they will visit more frequently and regularly leading to higher milk production. Forced cow traffic makes it possible to use forage at the bunk to provide motivation and while it reduces the number of fetch cows, it decreases the number of meals, decreases feed intake in some studies and decreases cow welfare because of increased standing time. Free cow traffic provides greater cow comfort and is preferred, but motivation to visit the robot is provided solely by the concentrate fed in the milking box. Hard, dust free pellets made of palatable ingredients such as barley and oats fed at a rate of 5 to 17 lbs per day result in the highest visit frequency and highest milk production. Limiting the energy density and starch level in the mixed ration fed at the bunk also increases the motivation provided by the concentrate. Current recommendations suggest feeding a partial mixed ration formulated for a production level 15 lbs below the mean of the group, combined with 5 to 17 lbs. of pelleted concentrate fed according to production in the robotic milking stall. While the need to use feed to stimulate milking visits creates additional challenges for the nutritionist and feed advisor, the ability to collect a great deal of data on the individual cow and to feed and supplement her individually also creates many new opportunities for more precise and individualized ration delivery.

<sup>1</sup>Contact at: 814471 Muir Line, Woodstock, Ontario, Canada, N4S 7V8, 519-467-5294, FAX 519-476-5845, Email:jack@dairylogix.com

**Introduction** – Robotic milking is being accepted as a viable alternative to milking parlors throughout the developed world. In areas such as Western Europe and Scandinavia where labor costs are high, and farms sizes are moderate, robotic systems have become the favored choice for new milking equipment purchasers. As labor costs and the complexity of labor management increase in North America and as the cost and challenges of managing robotic systems decline, robotic milking is earning its place as a viable option here as well. Currently an estimated 500 dairies in North America are milking with either the Lely Astronaut (Lely USA Inc., Pella, Iowa) or the DeLaval VMS (DeLaval Inc., Kansas City, Missouri) system, in herds ranging in size from 40 to 1200 cows. Both of these systems use a single milking stall, which the cow visits voluntarily. In both systems the primary motivation for cows to visit the stall is her desire and need to eat. The feed that attracts her can be in the concentrate feeder in the milking box, or with forced traffic, from this concentrate in combination with the feed at the bunk which is only accessible en route through or past the milking stall. Health concerns particularly problems with locomotion reduce visits to the milking stall and may be influenced by the diet. Greater milking frequency and more uniform milking times resulting from more visits to the milking stall increase milk production. Hence feeding management takes on a more complex and important role in the robotic dairy than it had with conventional parlor milking. A completely different approach to feeding is required. When nutrition advisors fail to recognize this, new robotic milking herd start ups often have very disappointing results.

**Goals in feeding the robotic dairy** -The goals of traditional dairy feeding programs include meeting the nutritional requirements of the cow in a way that ensures that she stays healthy, using feed ingredients that are economical and using labor efficient and cost effective feed delivery systems. With robotic milking there is a very important fifth goal: enticing the cow to visit the robotic milking stall regularly and frequently.

**The performance spiral when milking is voluntary** - Interactions among the activity or behavior of the dairy cow, her diet and feed consumption, her health and her milk production are complex and become even more complex with voluntary milking. Part of the complexity among these relationships is that none can claim to be distinctly “cause” and none is distinctly “effect”. For example, standard feeding management advice encourages producers to provide fresh feed more often, to stimulate a change in behavior, in the form of more frequent meals. This change in behavior is predicted to “cause” a change in diet, in the form of higher feed intake, which subsequently “causes” higher milk production. Alternatively diet may drive behavior, when a low fiber, high grain ration, is blamed for a high incidence of lameness, “causing” a change in behavior in the form of fewer trips to the feed bunk, subsequently “causing” lower feed intake and lower production. But when 3x milking, elicits an 8 to 12 lb. production response, higher production “causes” greater feed intake. In these examples each of the four attributes, behavior, diet, health and production is “cause” in some cases, and effect in others. When cows are milked at fixed intervals, external control of the “milking frequency” variable may limit variation in the other attributes. For example, under conditions of heat stress, cows reduce their activity and reduce their feed intake. Production suffers, but

twice daily milking, provides a baseline stimulus for production. Robotic milking is voluntary and variable, adding a new dimension to these interactions. If hot weather, reduces activity, it results in both lower feed intake and reduced milking frequency. Without a fixed, milking interval, heat stress in the robotic herd could start a downward spiral of reduced interest in feed, leading to less frequent milking, leading to lower production, and in turn even less interest in feed, etc. Based on this example, feeding management and an understanding of the interactions between diet, behavior, health and production take on a greater importance when robotic milking is considered.

**Using Feed as an Enticer for Robotic Milking** - Early research with robotic milking showed that without a feed incentive voluntary attendance at the milking stall is poor and highly variable. Feeding concentrate in the milking box, or forage or concentrate after passing through the milking box (forced cow traffic) improves attendance for milking. Although all commercial robotic systems currently offer concentrate in the milking box, and some use a form of forced cow traffic, failure of some cows to attend voluntarily remains a concern. The number of cows which must be fetched has been reported to be as low as 6% (Van't Land, 2000) on Dutch farms and as high as 19% on commercial farms in Ontario (Rodenburg and Wheeler, 2002). In recent years design improvements that have made the cow more comfortable in the milking stall, such as more space and the removal of the butt plate and adjustable manger in some models has improved milking frequency and reduced the number of cows fetched. In systems that still use these space limiting devices, adjusting them properly is an important factor in improving voluntary attendance.

A general understanding of eating behavior is useful for assessing how eating is altered by robotic milking feeding strategies. (Dado and Allen, 1994) reported that cows in a tiestall barn spent 300 minutes per day eating, 11 meals of 5.1 lbs for high producers and 3.7 lbs for low producers. These cows drank water 14 times per day, while cows in loose housing (Andersson, 1985) drank 6.6 times per day. In a freestall setting cows consumed 12.1 meals of TMR daily (Vasilatos, 1985). In contrast, a robotic herd with forced cow traffic, consumed 4.4 meals with a duration of 52.5 minutes and an average meal size of 8.8 Kg. (Tolle et.al., 2002)

**Forced vs. Free Cow Traffic** – Numerous studies have shown that attendance, while no longer “voluntary” in the pure sense, can be improved by forcing the cow to enter the robotic milking stall or an associated selection gate en route from the resting area to the feed manger or on her return from the manger to the resting area. This is commonly referred to as “forced” cow traffic. There are at least four common variations of “cow traffic” strategies used in robotic milking herds today. (1) Free cow traffic, where cows can access feeding and resting areas of the barn with no restriction. (2) Forced cow traffic with one way gates blocking the route from the resting area from the feeding area so cows leaving the resting area must enter the milking box, to be milked if the interval since the last milking makes her eligible, or “refused” if the milking interval is too short. After passing through the milking stall, the cow is released to the feeding area and can only return to the resting area through a one-way gate. (3) Forced cow traffic with “pre-selection” adds an entry lane where a sort gate directs cows eligible for milking to the holding area and ineligible cows to the feeding area. This reduces waiting times for

milking and for feed because only cows eligible for milking pass through the milking stall. Pre-selection can also be provided by selection gates in crossovers away from the robot, which open only for cows ineligible for milking. (4) Feed first forced traffic is a reversal of (2) which allows cows access to the manger from the resting area via one way gates, but they can only return to the resting area through the robotic milking stall, or through pre-selection gates that direct cows ineligible for milking directly to the free stalls or bedding pack.

Numerous studies report slightly higher milking frequency and a much-reduced need to fetch cows with forced traffic. (Hoogeveen, 1998; Van't Land, 2000). (Harms, 2002) reported 2.29, 2.63 and 2.56 milkings and 15.2, 3.8 and 4.3 fetching acts per day with 49 cows in free, forced and forced with pre-select traffic respectively. The number of meals was higher at 8.9 with free cow traffic, than with either forced or forced with pre-select, when cows consumed 6.6 and 7.4 meals respectively. Forage intake decreased when cows were switched to forced traffic and went back up in the forced with pre-select phase. (Thune, 2002) reported 1.98, 2.56 and 2.39 milkings, and 12.07, 3.86, and 6.46 feeding periods with free, forced and forced with pre-selection traffic respectively. On 7 Ontario farms with forced cow traffic (Rodenburg and Wheeler, 2002), average number of daily visits per cow, and therefore visits to the manger to consume TMR was  $3.40 \pm 0.44$ . This is many meals fewer than the 12.1 (Vasilatos, 1980) per day reported in a trial with free access and parlor milking. Fewer meals are associated with lower dry matter intake (Dado and Allan, 1994) and forced cow traffic has been shown to have this effect (Prescott et.al., 1998). Pre-selection systems result in some improvement in feed access but number of meals remains lower than with free traffic. Cows in forced traffic situation

also spend more time waiting for milking and less time lying down, (Winter and Hillerton, 1995). It is also of some concern that when a cow is in pain from a clinical case of mastitis or when she is lame, she will avoid milking in a free traffic situation and this alert the herdsman to her plight. Faced with the choice of starvation or milking this cow is more likely to go unnoticed in a forced traffic setting.

In the most recent comprehensive comparison for the two traffic systems (Bach et. al., 2009), cows were fed a partial mixed ration and up to 6.6 lbs of concentrate in the milking stall. Results summarized in table 1, illustrate that milking behavior, eating behavior and milk composition were all influenced by the choice of traffic system, but total dry matter intake and milk production were similar.

From a feeding standpoint forced traffic reduces the need to provide highly palatable feed in the robotic milking stall. As long as there is no alternative, most cows will go through the robotic milking stall out of sheer need to consume the ration at the feed manger, but reduced number of meals, reduced feed intake, reduced resting time, and longer waiting times, especially for timid cows make this system less desirable from the stand point of cow welfare and long term productivity.

With current technology there are numerous examples of robotic milking herds with free traffic that report over three milkings per day and very few fetch cows, and there are also numerous examples of forced traffic herds that report high feed intake, good production and few health issues. This demonstrates that both systems can work successfully under ideal circumstances. But when less than ideal conditions prevail, with free traffic the dairyman suffers the consequences in the form of fewer milkings and more fetch cows. With forced traffic the cows suffer the consequences with lower feed intake,

and longer waiting times. Since problems are much more likely to be resolved quickly when the dairyman suffers, free cow traffic is the preferable management system.

**Feeding Concentrate in the Milking Box** – Typical eating rate for pelleted concentrates is 0.45 to 0.65 lbs. per minute. Since cows spend 6 to 8 minutes in the stall per milking maximum concentrate fed during milking is 2.5 to 3.5 lbs, or 7.5 to 10.5 lbs per day for a cow visiting three times. Some herds are successfully increasing pellet delivery rates in the robot to as much as 1 lb per minute and 18 lbs per day without seeing feed left behind. Additional grain is usually fed as part of a mixed ration in the manger or in individual feeders in the barn. The use of computer feeders with robotic milking can be organized strategically so that cows that require additional concentrate can receive it in computer feeders. These feeders can be linked to the milking software so that cows can only use them while they are ineligible for milking, or they can be located in a special exit area from the robot to further encourage traffic to the milking stall.

The concentrate fed in the milking stall is the “candy” that attracts the cow to come to the stall frequently for milking. More frequent milking shortens milking intervals and decreases variation in milking interval. Both of these outcomes increase milk production. Having fewer cows to fetch reduces labor for the operator.

The importance of feeding palatable concentrate in the milking stall, is illustrated by a case study on one Ontario farm. (Rodenburg and Wheeler, 2002) Initially, a low cost pellet formulated with lower palatability ingredients including gluten meal, canola and tallow was fed. Poor pellet strength caused a build up of fines in the bottom of the feeders. A stronger pellet of high palatability containing 3 (vs. 0) % molasses and 96 (vs.



65) % high palatability ingredients was substituted. Voluntary visits increased from 3.40 to 4.04, and voluntary milkings from 1.72 to 2.06 per cow per day. Canadian robotic milking system owners describe cows that they have to fetch for milking, as “lazy” when there is no clear reason, such as inexperience, clinical mastitis or lameness, for not attending voluntarily. Using this definition, “lazy milkings” and “lazy cows” declined from 27.3% and 16.0% to 12.7% and 7.1% respectively, when the stronger pellet replaced the weaker one. In another study (Rodenburg, Focker and Hand, 2004) we formulated a concentrate for what we thought was maximum palatability. Ingredients included corn, soya hulls, wheat shorts, barley, bakery meal, soybean meal, corn distillers, extruded soy meal, wet molasses, animal vegetable fat blend, vitamin mineral premix, sodium bicarbonate, salt, pellet binder and fenugreek flavour. In comparisons to commercial concentrates on four farms, in trials with three consecutive 15-day crossover/switchback feeding periods we found that visits (3.95 vs. 4.80) and milkings (2.69 vs. 2.81) were fewer ( $p < .05$ ) for the experimental pellet when compared to a stronger commercial pellet (shear strength of 91.2 vs. 96.0 pdi) in trial 1. In trial 2, the experimental pellet was compared to a different commercial product of equal shear strength and in this trial attendance was unaffected. In trial 3, conducted in the same herd as trial 2, the pellet was reformulated to exclude all mineral ingredients, but no difference in attendance was found. In trial 4 a mixture of 50% commercial pellets and 50% high moisture corn was compared to our experimental pellet, adjusted to make it isonitrogenous with the control. As in trial 1, number of visits (3.06 vs. 3.33) and milkings (2.34 vs. 2.49) were lower ( $p < 0.05$ ) for the experimental pellet. In this trial shear strength of the experimental pellet was weaker, 86.9 pdi vs. 97.7 pdi, than the

commercial pellet and there was evidence of fines in the feeder when it was fed. One other herds volunteered to test a mixture of 49 % dried corn distillers, 49% cracked corn, 2% molasses and 0.1% flavoring agent fed in a mash form, but during over a 6 day feeding period the number of visits decreased from 3.93 to 3.57, and number of milkings from 2.50 to 2.35. Milk production declined from 57.2 to 53.6 lbs and the trial was discontinued. These studies clearly demonstrate that the concentrate fed in the robot should be pelleted and pellets should be of high quality and free from fines. Feed delivery systems should be designed to minimize pellet breakdown during handling.

More recently Danish researchers (Madsen et. al., 2010) compared 7 pellet formulations and found substantial differences in the number of visits, the number of milkings, the number of fetch cows and in milk production, linked to the ingredients used in the pelleted concentrates. Results are summarized in table 2. As illustrated cows preferred a barley and oats combination, followed by a wheat based concentrate. Corn was less palatable and a fat enriched pellet and one based on dried grass resulted in significantly fewer visits and lower milk production. Danish workers have also demonstrated a preference for barley oats mixtures over corn in other studies with computer feeders. In studies of feed palatability higher intake for flavored concentrates (Arave, 1989) and sweeteners (Weller, 1989; Nombekala, 1994) has been reported in some trials, but not in others (Murphy, 1997). Published palatability ratings for feed ingredients tend to be based on field experience rather than controlled studies. (Amaral-Phillips, 1993; Maiga, 1997) Highest palatability is assigned to brewers grains, distillers grains, hominy, molasses and beet pulp. Soybean meal, roasted soybeans, corn, barley and wheat midlings rank intermediate; raw soybeans, and canola meal are rank low, and

corn gluten meal, blood, meat and fish meals, tallow, bypass fats, mineral mixes, buffers and niacin rank very low. Pellets are clearly favored over mash, and heat treated rapeseed meal, barley with 10% rapeseed fatty acids, or with 10% palm oil, or with 10% glycerol were all preferred over ground palm kernel expeller (Sporndly and Asberg, 2006).

DeLaval robotic specialists suggest that caramel flavoring added at 0.5 lbs per ton to the robot pellet is thought to enhance palatability (Futcher, 2011)

The amount of pellets fed in the robotic milking stall appears to have less influence on visiting behavior than the composition and pellet strength. Feeding 6.6 or 17.6 lbs of pellets in the robotic milking stall to cows fed a high corn silage diet at the manger did not result in any difference in the number of milkings or the number of cows that required fetching. (Bach et. al. 2007). In this study the ration fed was quite energy dense, and it is likely that this reduced the attraction offered by higher levels of concentrate in the milking stall. A study at the University of Ghent (Hauspie, 2008) summarized in table 3 found that visits and milk production increased when the amount of concentrate in the mixed ration in the manger was reduced by 30% or 4.0 lbs. per cow and the amount of concentrate fed in the milking stall was increased by 12% or 1.54 lbs. Despite lower grain feeding, milk production went up in response to more frequent milking.

Varying the amount of concentrate fed in the milking stall according to production can also decrease grain feeding and associated feed cost. Since the concentrate dispenser in the robotic milking stall delivers feed on a volume basis, it is essential that it be calibrated after each new load of feed is delivered and on a regular basis in between deliveries as well. Pellet ingredients, pellet strength and quantity fed can all affect visit behavior, and visits and milkings drive production, so nutritionists need to pay careful

attention to pellet formulation, manufacturing and handling. This product is the “candy” for robotic milking units.

**Grazing and Automatic Milking** - Grazing and automatic milking have been successfully combined in research studies (Sporndly and Wredle, 2002; Ketelaar-DeLauwere et.al., 2000) and on commercial farms (Jagtenberg and VanLent, 2000). When distances to pasture increase, especially beyond 400 meters (Wiktorsson and Sporndly, 2002; Sporndly and Wredle, 2004) milk production, milking frequency and grazing time for late summer pasture declined suggesting pastures close to the barn are preferred. Providing more supplementary forage in the barn did not increase milk yield in cows grazed 260 meters from the robotic milking system. (Sporndly and Wredle, 2004). With distances to pasture of up to 330 meters, no differences in milking frequency or milk production were found when water was offered only in the barn vs. in the barn and in the pasture. (Sporndly and Wredle, 2005)

Traffic management strategies that offer water and supplementary feed prior to milking and direct cows back to pasture are suggested as a way to encourage frequent attendance. The Lely company recommends strip grazing and directing cows to new pastures twice daily using a selection gate at the barn exit that directs cows ineligible for milking to a new pasture twice daily. (Van Mourik et.al., 2010)

**High Grain Diets and Automatic Milking** - In North America, the more continental climate favours diets of whole plant corn and alfalfa silage, which are more easily fed from storage. Because grain, is inexpensive and investment in housing is high, diets with

a high concentration of grain, which support high milk production are favoured. Mixing all grain, forage and supplement ingredients into a total mixed ration provides the accuracy of formulation and control of fibre level needed to minimise the risk of digestive disturbance with these diets. The need for concentrate in the milking box, conflicts with traditional feeding practices in the US and Canada. Purchased pelleted concentrates cost more than the high moisture grains they replace. Feeding grain separate from the TMR can lead to situations where the maximum concentrate to forage ratio is exceeded when TMR intake is depressed. Field data (Rodenburg and Wheeler, 2002) suggests that in robotic milking herds where a high grain TMR is fed at the manger, frequency of voluntary milking is lower and more cows must be fetched as illustrated in Fig 1. Measures of voluntary milking appear impaired in diets with more than 1.66 Mcal Nel per kilogram dry matter or more than 48% concentrate.

High grain diets are associated with laminitis (Manson, 1988a), and perhaps the three farms above 1.66 Mcal Nel/Kg in Fig. 1 suffer from a level of “subclinical” laminitis, which is decreasing the mobility of cows. Carbohydrate level and fermentation rates, matching rumen availability of protein, and the level and form of dietary fiber are key factors which influence rumen acidosis and laminitis. Limits of 25 to 35 % NDF, with 75% from forage, 35 to 40% NSC, 30 to 40% starch in the diet dry matter, and a ratio of forage NDF to ruminally degradable starch of > 1:1 have been recommended. (Nocek, 1997).

An Israeli trial assessing the impact of replacing pellets made with high starch grains with isocaloric pellets made with soy hulls and corn gluten high in digestible fiber

(Halachmi et. al., 2009) reported higher milk production (94 vs. 86 lbs) but milking frequency at 3.12 and 3.16 visits per day was not different.

According to NRC predicted dry matter intake for a cow fresh 11 days producing 90 lbs. of 3.5% fat, 3.0% protein milk is 36.8 lbs. or 35% less than for the same production at 90 days. (NRC, 2001). When a portion of the grain is fed separately in the milking box, TMR intake depression in early lactation means that a small amount of grain can cause the above guidelines to be exceeded. Grain fed in the milking box should be limited to 4.4 lbs. per day at calving and increased slowly over several weeks while appetite and TMR intake stabilize.

Cows on high grain diets may also be less aggressive due to a direct metabolic effect. Cows on high grain diets spend less time eating and ruminating and more time resting (Robinson, 1997), and consume fewer meals (Friggens, 1998). With fewer meals, directed cow traffic becomes less effective.

The type of diets described as high grain and high energy in this paper are typical of programs commonly used with high producing TMR herds in North America. If these diets result in poorer voluntary attendance for milking and lower milking frequency, understanding this relationship better will be an important area of future research.

The current trend in AMS herds in Canada is toward less grain feeding. The standard recommendation from Lely (VanMourik et. al., 2008b) is to balance the partial mixed ration (PMR) fed in the manger for a production level that is 15 lbs below the average production of the group, and to selectively feed pelleted concentrate at a rate of 5 to 17 lbs. per day according to production for all cows fresh more than 4 weeks. When a new herd is started on robotic milking cows should be acclimatized to the pellets by

feeding them at the manger until the robotic milking stalls are installed. Switch the concentrate to milking stall at start up. DeLaval recommends a default value of 8 lbs of concentrate per cow in the milking stall at start up, combined with a mixed ration in the manger. (Futcher, 2011) After start up concentrate feeding can be adjusted using tables with a range of 4 to 14 lbs per cow per day with free traffic. More moderate upper limits are suggested for forced traffic applications. Grain feeding in the first two to three weeks of lactation is very conservative. In the past it was common for rations to be formulated to provide 110 to 115% of the nutrient requirements of the average cow. These diets are now being evaluated for “feeding efficiency” which the feed advisor defines as the measure of how closely the feed provided supplies the nutrients required at the current production level. Response to this shift in emphasis appears to be positive as producers report higher forage intakes, steady production and higher attendance. In many herds, the stimulus of increased milking frequency has increased milk production, despite lower levels of grain feeding.

### **Dynamic Feeding and Other opportunities for Greater Feeding Precision -**

Currently, software can be purchased for Lely robotic milking systems that automatically optimizes the robot grain allocation for each cow in the herd based on feed and milk prices, and yesterday’s production, milk composition and milking speed of each cow, using a dynamic linear model (VanHolder et.al 2010). Although the concept of adaptive feeding models that base today’s feed allocation on how the cow has responded to feeding changes in the recent past is a valid one, the present model has not been verified under North American conditions. Farmers in Europe report that during the period of low

milk prices in 2009, dynamic feeding reduced grain feeding levels for many cows and increased income over feed costs. (Wesselink, 2011).

The concept of dynamic feeding illustrates that robotic milking provides a unique opportunity to feed cows individually. The evolution of total mixed rations over the last 40 years has meant that the concept of feeding the individual cow according to her nutrient requirements as well as her individual behavior and preferences has fallen by the wayside. Robotic milking systems are available with the capability to feed several feed types in pellet, mash or liquid form. These systems provide daily data on milk production, milk composition, milking and eating behavior and a wide variety of other parameters. Add on precision management tools such as DeLaval's Herd Navigator in line testing for milk urea nitrogen, and beta hydroxy buterate will add further information to enhance individual cow nutrition management. While the need to attract the cow to the stall with feed creates additional challenges for the nutritionist and feed advisor, the capability for gathering detailed information about individual cows and the capability to provide a wide variety of feeds and additives on an individual basis creates many new opportunities as well.

## **References**

Amaral-Phillips, D. M., and R. W. Hemken. 1993. Using by-products to feed dairy cattle. University of Kentucky. On-line publ. ASC-136.

Andersson, M. 1987. Effects of Number and Location of Water Bowls and Social Rank on Drinking Behaviour and Performance in Loose Housed Dairy Cows. *Livestock Production Science* 17:19-31

Arave, C. W., D. Purcell, and M. Engstrom. (1989) Effects of feed flavors on improving choice of ten percent meat and bone meal dairy concentrate. *J. Dairy Sci.* 72:563.



Bach, A., C. Iglesias, S. Calsamiglia, and M. Devant, (2007) Effect of Amount of Concentrate offered in Automatic Milking Systems on Milking Frequency, Feeding Behavior, and Milk Production of Dairy cattle Consuming High amounts of Corn Silage, *Journal of dairy Science* 90-11, pg 5049-5055

Dado, R. G. and M. S. Allen. (1994) Variation in and relationships among feeding, chewing, and drinking variables for lactating dairy cows. *J. Dairy Sci.* 77:132

Futcher, M (2011), DeLaval Marketing Manager for Automatic Milking.  
[Mark.futcher@delaval.com](mailto:Mark.futcher@delaval.com), personal communication

Friggens, N. C, G. C. Emmans, I. Kyriazakis, J. D. Oldham, and M. Lewis. (1998) Feed intake relative to stage of lactation for dairy cows consuming total mixed diets with a high or low ratio of concentrate to forage. *J. Dairy Sci.* 81:2228.

Harms, J., G. Wendl, and H Schon, 2002. "Influence of Cow Traffic on Milking and Animal Behaviour in a Robotic Milking System" in Proceedings of the First North American Conference on Robotic Milking, March 20-22, 2002, Toronto Canada, Wageningen Press, Pp II 9-II 14

Hauspie, 2008. Reported in the Lely Monthly Management Magazine, July 2008, Lely Industries, Weverskade 10, 3155 PD, Maasland, The Netherlands

Hogeveen, H., A.J.H. van Lent, and C.J. Jagtenberg 1998. "Free and One-Way Cow Traffic in Combination with Automated Milking" Proceedings of the 4<sup>th</sup> International Dairy Housing Conference St. Louis Missouri January 28-30 1998 ASAE pp. 80-87

Jagtenberg, K., and J. Vincent (2000) Robotic Milking and Pasturing. *Praktijkonderzoek* 13 (3) 7-9

Keetelaar-De Lauwere, C.C., A.H. Ipema, C. Lokhorst, J.H.M. Metz, J. Noordhuizen, W.G.P. Schouten, A.C. Smits, (2000) Effect of Sward Height and Distance Between pasture and Barn on Cows Visits to an Automatic Milking System and Other Behaviour. *Livestock production Science* 65(1-2): 131-142

Madsen, J., M.R. Weisbjerg, and T. Hvelplund, 2010, Concentrate composition for Automatic Milking Systems- Effect on Milking Frequency. *Livestock Science* 127, 45-50

Manson, F.J. and J.D. Leaver, 1988a, "The influence of concentrate amount on locomotion and clinical lameness in dairy cattle." *Anim. Prod.* 47:185

Maiga, H.A., G.D. Marx, V. W. Crary and J. G. Linn. 1997. Alternative Feeds for Dairy Cattle in Northwest Minnesota: An Update. *Dairy Update Issue* 126.

[www.ansci.umn.edu/dairy/dairyupdates/du126.htm](http://www.ansci.umn.edu/dairy/dairyupdates/du126.htm)

Murphy, M.R., A.W.P. Geijssel, E.C. Hall, and R.D. Shanks, 1997, "Dietary variety via sweetening and voluntary intake of lactating cows" *J. Dairy Sc.* 80:894-897

National Research Council, 2001, *Nutrient Requirements of Dairy Cattle*, Seventh Revised Edition, National Academy Press, Washington D.C.

Nombekela, S. W., M. R. Murphy, H. W. Gonyou, and J. I. Marden. (1994) Dietary preferences in early lactation cows as affected by primary tastes and some common feed flavors. *J. Dairy Sci.* 77:2393.

Robinson, P. H. and R. E. McQueen. (1994) Influence of supplemental protein source and feeding frequency on rumen fermentation and performance in dairy cows. *J. Dairy Sci.* 77:1340.

Prescott, N.B., T.T. Mottram, and A.J.F. Webster, 1998, Relative motivations of dairy cows to be milked or fed in a Y- maze and an automatic milking system. *Applied Animal Behaviour Science* 57, 23 -33

Rodenburg, J. and B. Wheeler, 2002, Strategies for Incorporating Robotic Milking into North American Herd Management. In: *Proceedings of the first North American Conference on Robotic Milking*, Toronto, Canada, pp III 18 – III 32

Rodenburg, J. E. Focker, K. Hand, 2004, Effect of the Composition of Concentrate Fed in the Milking Box, on Milking Frequency and Voluntary Attendance in Automatic Milking Systems, *Proceedings of the International Conference on Automatic Milking*, Lelystad, March 2004, pg 511

Sporndly, E., E. Wredle, 2002, The Effect of Distance to pasture, and Level of Supplementary Feeding on Visiting Frequency, Milk Production and Live weight of Cows in an Automatic Milking System, *Proceedings of the First North American Conference on Robotic Milking*, Toronto Canada, March 2002, III-76-77

Sporndly, E., and E. Wredle, 2004, Automatic Milking and Grazing – Effects of Distance to Pasture and Level of Supplements on Milk Yield and Cow Behaviour. *Journal of Dairy Science* 87 -6, pg 1702-1712

Sporndly, E., and E. Wredle, 2005, Automatic Milking and Grazing – Effects of Location of Drinking Water on Water Intake, Milk Yield and Cow Behaviour. *Journal of dairy science* 68-5, pg 1711-1722

Sporndly, E., and T. Asberg, 2006, Eating Rate and Preference of Different Concentrate Components for Cattle, *Journal of Dairy Science*, 89-6 pg 2188-2199

Tolkamp, B. J., D. P. Schweitzer, and I. Kyriazakis. (2000) The biologically relevant unit for the analysis of short-term feeding behavior of dairy cows. *J. Dairy Sci.* 83:2057.

Tolle, K.H. O. Joulaud, G. Janknecht, G.Krieter, 2002. Feed Intake of Dairy Cows in Automatic Milking Systems with Forced Cow Traffic. *Zuchtungskunde* 74 (5): 330-40

Thune, R.O., A.M. Berggren, L. Gravas, and H. Wiktorsson, 2002, Barn Layout and Cow Traffic to Optimize the Capacity of n Automatic Milking System.” in Proceedings of the First North American Conference on Robotic Milking, March 20-22, 2002, Toronto Canada, Wageningen Press, Pp II 45-II 50

VanHolder, T., P. Kool, and P.P.J. van der Tol, 2010, Health and Production Characteristics for a Dynamic feeding Model, in proceedings of the First North American Conference on Precision Dairy Management, March 2-5, 2010, Toronto, Canada, [www.precisiondairy2010.com](http://www.precisiondairy2010.com) pg 206-207

Van Mourik, J.D., C.J. Hollander, and A. Hempenius, 2008, Lely Monthly Management Magazine, May 2008, Lely Industries, Weverskade 10, 3155 PD, Maasland, The Netherlands

Van Mourik, J.D., C.J. Hollander, and A. Hempenius, 2008b, Lely Monthly Management Magazine, July 2008, Lely Industries, Weverskade 10, 3155 PD, Maasland, The Netherlands

Van'tLand, A., A.C. Van Lenteren, E. Van Scooten, C. Bouwmans, D.J. Gravesteyn and P. Hink, 2000, Effects of Husbandry System on the Efficiency and Optimization of Robotic Milking Performance and Management. In *Robotic Milking: Proc. of the International Symposium held in Lelystad, the Netherlands 17-19 August 2000*, Wageningen Press. pp 167-176

Vasilatos, R. and P. J. Wangsness. (1980) Feeding behavior of lactating dairy cows as measured by time-lapse photography. *J. Dairy Sci.* 63:412.

Weller, R.F., and R.H. Phipps, 1989, Preliminary studies on the affect of flavouring agents on the dry matter intake of silage by lactating dairy cows” *J. agric. Sci. Camb.* 112:67-71

Wesselink, W., 2011, Dynamic feeding, in *Western Dairy Farmer* magazine, January/February 2011 pg 20-26

Wiktorsson, H, E. Spordly, 2002, Grazing: An Animal Welfare Issue for Automatic Milking Farms, In: *Proceedings of the first North American Conference on Robotic Milking*, Toronto, Canada, pp VI 32 – 42

Winter, A., and J.E. Hillerton, 1995, Behaviour associated with feeding and milking of early lactation cows housed in an experimental automatic milking system. *Applied Animal Behaviour Science* 46, 1-15

**Table 1: (Bach et. al. 2009) Feeding and milking behavior, and milk production and composition of cows with free vs. forced traffic.**

Item (per cow per day)	Treatment		SE	P-value
	<u>Free traffic</u>	<u>Forced traffic</u>		
Total milkings	2.2	2.5	0,04	<0.001
Fetches milkings	0.5	0.1	0.03	<0.001
Voluntary milkings	1.7	2.4	0.06	<0.001
PMR intake (lbs. DM)	41.0	38.8	1.34	0.24
No. of meals of PMR	10.1	6.6	0.30	<0.001
Concentrate intake (lbs.)	5.5	5.5	0.09	0.99
Milk production (lbs.)	65.7	68.1	1.74	0.32
Milk fat content (%)	3.65	3.44	0.078	0.06
Milk protein content (%)	3.38	3.31	0.022	0.05

**Table 2. (Madsen et. al.) Effect of concentrate formulation on robotic milking behavior and milk production.**

Concentrate (per cow/ day)	Standard (Mean)	Barley	Wheat	Barley/Oats	Corn	Fat Rich	Dried Grass
		(Effect of test feed expressed as test feed minus standard)					
Milkings	2.96	-0.03	0.17	0.35**	0.02	-0.36*	-0.93***
Refusals	2.09	-0.05	0.44	1.87	0.31	-0.39	-1.16
Fetchings	0.026	0.028	0.019	0.009	0.50	0.042	0.17
Lbs.Milk	57.5	0.22	3.53*	2.65	0.44	-1.98	-9.04***

\*\* = p < 0.01, \*\*\* = p < 0.001

**Table 3. (Hauspie, 2008) Milking frequency and milk production response to feeding more of the concentrate in the robotic milking stall**

(Per cow per day)	<u>Control</u>	<u>Treatment</u>	<u>%Difference</u>
Concentrate in the manger (lbs.)	13.2	9.3	- 30%
Concentrate in the milking stall (lbs.)	12.8	14.3	+ 12%
Milkings	2.3	2.5	+ 8%
Refusals	1.0	1.4	+27%
Milk production (lbs.)	60.4	65.0	+ 7%

**Fig. 1 (Rodenburg and Wheeler, 2002) Energy Level in Diet Dry Matter and Milking Behavior .**

